



SEDIMENTOLOGY OF EGYPT

ISSN 1110-2527


JOURNAL OF THE SEDIMENTOLOGICAL SOCIETY OF EGYPT

Volume 14

CONTENTS

January 2006

Coastal Sediments and Their Polluting Metals of El-Hamrawein Harbour, Egyptian Red Sea Coast: Chocs for Monitoring Environmental Hazards. Hashem A. Madkour, Abn El-hagag N. Ahmed and Ahmed W. Mohamed.....	1
Hydrodynamic Factors Affecting Sedimentation Regime in Damietta Harbour, Nile Delta Coast, Egypt. Abo Baker I. Abo Zed.....	15
Beach Changes Along the Eastern Nile Delta Coast and Its Relation with the Coastal Processes and the Constructed Detached Breakwaters, Egypt. Mohamed Abdel Gull.....	29
The Effect of Engineering Structures on the Morphology and Sedimentology of Ras El Barr Beach Magda M. Abo El-Safa.....	41
Dynamic Processes Affecting Marakia and Marina Beaches Northwestern Coast of Egypt. Abo Baker I. Abo Zed, Kadry N. Sediek, samy M. Shereet, and Abdel Moneim A. Bndr.....	53
Mineralogy and Sedimentary Studies on Recent Coastal Sabkha, Northwestern Mediterranean Coast, Egypt. Shams El-Din M. Shaheen.....	65
Hydrochemical and Isotopic Characteristics of the Groundwater in Central and South Eastern Desert, Egypt. Sawssan G. Abd El Samie, Ewaia M. Monssa, Mohamed A. Sayed.....	83
Hydrostratigraphy of Central Nile Delta (Egypt) Using Geoelectric Measurements Mohamed G. Atwia, Mohamed M. Abon Heleika, Ahmed M. El-Shishtawy, and John M. Sharp Jr.	95
Miocene Rocks and Their Impact on Shallow Groundwater, Siwa Oasis, Western Desert, Egypt. Bothainn M. Monssa.....	117
Contributions to the Hydrochemical Characteristics and Some Hydrogeological Aspects of the Different Groundwater Aquifers in Siwa Oasis, North Western Desert, Egypt. Rafik G. Fathy, Abd El Naem H. Afifi and Mohamed El-Amir Nagaty	143
Sedimentary Facies Characteristics, Facies Development and Sequence Stratigraphic Implications of the Jurassic Shusha Sediments in Gabal El-maghara, North Sinai, Egypt. Monir H. El-Azabi.....	161
Pervasive Dolomitization of the Cenomanian Sequence in the Gabal Manzour Area, North Sinai, Egypt. Rifai I., Rifai and Ahmed S., Mansour.....	199
Role of Epigenetic Processes in the Enrichment of Mineralized Dolostones of Um Bugna Formation, Southwest Sinai. Ahmed E. Shata.....	219
Minerals Controlling the Radioactivity in the Stream Sediments at Wadi El Shaffal Area, Southwest Sinai, Egypt. Salah S. El-bulakssy.....	243
Sedimentology, Mineralogy and Geochemistry of the Early Paleozoic Paleoplacers in Both Eastern and Western Sides of the Gulf of Suez. Nasser L. El Agamri and Ahmed E. Shata	261
Carbonate Cementation Patterns in a Mixed Siliclastic / Carbonate Sequence: Evidence from the Lower Cretaceous (Fayed Formation), Southern Ismailia, Egypt. Alaa M. Salem, Ahmed M. El-Shishtawy, and Sadoon Morad.....	281
Contribution to the Geochemical Characteristics of Carbonaceous Shale, Qatrani Area, Western Desert, Egypt. Ewaia M. Monssa, and Yehia S. Haroun.....	295
Provenance and Palaeoenvironmental Conditions of Pliocene Sediments in the Area East of Higaza, Nile Valley, Upper Egypt. Mohamed R. Osman and Hanz Kurzweil.....	307
The Relationship Between the Quaternary Reef Terraces and the Beachrocks, Red Sea Coast Egypt: Sedimentological and Diagenetic Aspects. Essam M. El Khoriby and Tarek I. Anan.....	331
Invited Talk.....	361



Invited Talk

Mortada Mourad Taha El Aref

**Professor of "ore geology", Geology
Department, Cairo University**

EMPLOYMENT

- 1969-1975... Assistant, Geology Department, Faculty of Science, Assiut University
1975-1976... Assistant Lecturer, Geology Department, Faculty of Science, Assiut University
1976-1981... Ph D. Candidate, Heidelberg University, West Germany
1981-1986... Lecturer, Geology Department, Cairo University
1986-1993... Associate Professor (Ore Deposits), Geology Department, Cairo University
1993-Present... Professor of Ore Geology (Ore in Sediments), Geology Department, Cairo University
2003-present... Professor of Ore Geology (Ore in Sediments), Geology Department, Cairo University
Director of Cairo University Centre for Environmental Hazard Mitigation (CEHM)
Director of Regional Centre for Training and Technology Transfer for the Arab States
(BCRC- Cairo)

Egyptian Karst Morphology and Processes, Its Economic Potentiality and Environmental Impacts

Karst is internationally used term, originally the German from the Slavic word "kras" or "krs" meaning a bleak waterless place. The term describes a terrain generally underlain by limestone or dolomite, in which the topography is chiefly formed by the dissolving of rock, and which may be characterized by sinkholes, sinking streams, closed depressions, subterranean drainage and caves. In the temperate and tropical areas, where carbonate rocks are exposed, environmental water may combine with CO₂ in the atmosphere or in the vegetation cover above the rock to produce a weak solution of carbonic acid (H₂CO₃) which slowly dissolves the carbonate rocks.

The Egyptian Phanerozoic carbonate successions are punctuated and truncated by several paleokarst (unconformity) surfaces of local or regional magnitudes. The drowned (fossilized or buried) paleokarst surfaces include: a) Depositional paleokarst surfaces terminating meter scale shoaling cycles, generally with topographic relief of centimeters to decimeters. The effects of exposures are usually restricted to surface solution sculptures, near surface cementation and minor subsurface dissolution, and

b) Drowned paleokarst surfaces representing breaks in sedimentation during periods of relatively long-lived exposure and sea level fall as well as intensive karstification inducing lateritization and pedogenesis along paleohighs of regional or local scales. The associated karst features include widened karren, sinkholes, small scale dolines, collapse breccia, subsurface solution channels and caves together with residual soil products. Different types of enriched ore deposits are genetically related to paleokarstification and developed along paleokarst surfaces as they constitute substantial horizons of the paleokarst system and the resulted paleokarst profile. Among these karst-related deposits are:

1. Stratabound Carboniferous Mn deposits with Cu and U sulphates, phosphates, vanadates and carbonates, Cu chlorides, gibbsite, kaolinite and alunite. Um Bogma region, Sinai.
2. Stratabound Eocene Fe and Ba deposits, El Bahariya Depression, Western Desert.
3. Stratabound Miocene (post-Miocene ?) Pb, Zn, Sr and Ba deposits, Red Sea coastal zone, and
4. Karst calcareous deposits ("Egyptian Alabaster") mixed with terra rossa sediments and formed at the expense of different carbonate sequences along different paleoerosion surfaces.

The exposed Phanerozoic carbonates also show enormously varied surface and subsurface solution forms associated with surficial duricrusts and paleosols as a result of uplifting and long exposure. The resulted karst landforms are: a) Mature and degraded conekarst, b) Mature and degraded towerkarst, and c) flat-topped tablelands. Solution depressions of variable dimensions (sinkholes, dolines and poljes) and morphogenetic karren forms are widely distributed in the plain surfaces. Farafra depression is one of large poljes, formed by the coalescence of solution dolines and degradation of towerkarst system. Partially to completely filled small and large-scale subsurface cavern levels, potholes and channels are commonly developed along vertical joints and bedding planes.

The analyses of these features and the associated sediments indicate multi-cycles of karstification involving continuous lowering of the landscape. Paleotopography, paleoclimate, soilification, lithology of the host rocks and the tectonic evolution and related structural elements, are the main fundamental factors that controlled the formation, distribution and evolution of the karst landforms and the composition and textures of the associated karst products.

The environmental impacts of karst are either positive or negative. Among the positive impacts are: a) The beautiful landscape and natural surface and subsurface features, b) The economic potentiality of the karst products and precipitates, and c) The karst features and sediments are good indications of the paleoclimate, paleotopography and paleogeographic distribution of the paleohighs and the paleoshorelines. Among the negative impacts of karst are: a) Contamination of the groundwater, b) Household waste problems, c) Flooding, and d) Engineering problems (sinkhole collapse).



المجلة المصرية لعلم الرسوبيات

(مجلة الجمعية المصرية لعلم الرسوبيات)

ISSN 1110-2527

مجلد ١٤



يناير ٢٠٠٦

